



MASENO UNIVERSITY
UNIVERSITY EXAMINATIONS 2016/2017

**FIRST YEAR FIRST SEMESTER EXAMINATION FOR DEGREE
OF MASTER OF SCIENCE IN APPLIED STATISTICS**

CITY CAMPUS - REGULAR

MAS 809: EPIDEMIC MODELLING

Date: 2nd December, 2016

Time: 2.00 - 5.00pm

INSTRUCTIONS:

- Answer ANY THREE Questions.
- Observe further instructions on the answer booklet.

QUESTION ONE

[20 Marks]

(a) Explain clearly what is meant by the following;

- i. Epidemic
- ii. Endemic
- iii. An Epidemic model

(6 Marks)

(b) Consider the SI model defined by the following equations;

$$\frac{dS(t)}{dt} = -\lambda(t)cS(t)$$

$$\frac{dI(t)}{dt} = \lambda(t)cS(t) - vI(t)$$

where $\lambda(t) = \frac{\beta I(t)}{N(t)}$, $N(t) = I(t) + S(t)$ and c and v are some constants. Show that if $S(t) \cong N(t)$, then

$$I(t) = I_0 e^{(\beta c - v)t}$$

and the time taken, if there is an epidemic, for the number of infected to double is

$$t_d = \frac{\ln 2}{(\beta c - v)}$$

(10 Marks)

(c) Using the results in question (b), or otherwise, determine the conditions under which an epidemic will arise.

(4 Marks)

QUESTION THREE

[20 Marks]

(a) Briefly explain the main features of each of the following classes of epidemic models;

- i. Empirical models
- ii. Deterministic models
- iii. Stochastic Models

(10 Marks)

(b) Consider a household of size 7 with 2 zero generation cases;

- i. List all the possible epidemic chains.

(5 Marks)

- ii. Determine the probability that the size of the epidemic chain is five.

(5 Marks)

QUESTION FOUR

[20 Marks]

(a) Differentiate between the fixed effects and the random effects chain Binomial Models applicable in epidemic modeling. (10 marks)

(b) Consider the epidemic Chain Binomial model.

Let $i_t : t = 0, 1, \dots, r$, ($i_{r+1} = 0$), denote the number of infectives at time t since the onset of the epidemic, and $S(t) : t = 0, 1, \dots, r$, ($S_r = S_{r+1}$) be the number of corresponding susceptibles at time t .

Let q_i be the probability that a susceptible escapes infection when exposed to i infectives.

Determine Prob [epidemic chain is $i_0 \rightarrow i_1 \rightarrow \dots \rightarrow i_r$] and deduce the expression for the probability assuming;

- i. Reed Frost model
- ii. Greenwood Model

(10 Marks)

QUESTION FIVE

[20 Marks]

Consider the “general epidemic” model with susceptibles $S(t)$, infectives $I(t)$, and Removals $R(t)$. Assuming homogeneous mixing among these classes and with continuous time t , let

$$\begin{aligned}dS(t)dt &= -\beta S(t)I(t) \\ \frac{dI(t)}{dt} &= \beta S(t)I(t) - \alpha I(t) \\ \frac{dR(t)}{dt} &= \alpha I(t)\end{aligned}$$

with initial conditions $S(0), I(0), R(0) = (S_0, I_0, 0)$ and α and β are some constants. Also let $N(t) = R(t) + S(t) + I(t)$.

(a) Explain this formulation with respect to the spread of diseases.

(6 Marks)

(b) Show that,

i. $\frac{dI(t)}{dt} = \beta I(t)(S - \rho), \rho = \frac{\alpha}{\beta}$.

(4 Marks)

ii. $S(t) = S_0 e^{-\frac{R(t)}{\rho}}$.

(4 Marks)

iii. For $N(t)$ such that $N(t) = \rho + v$ where $v \ll \rho$ with $N(t) \cong S_0$;

$$R(\infty) = 2v$$

$$S(\infty) = \rho - v$$

Comment on your results.

(6 Marks)